



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 205/08	A1	(11) International Publication Number: WO 97/16424 (43) International Publication Date: 9 May 1997 (09.05.97)
(21) International Application Number: PCT/US96/17083 (22) International Filing Date: 30 October 1996 (30.10.96) (30) Priority Data: 60/006,182 2 November 1995 (02.11.95) US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). (72) Inventor: SHANKAR, Bandarpalle, B.; 1124 Van Arsdale Drive, Branchburg, NJ 08853 (US). (74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROCESS FOR PREPARING 1-(4-FLUOROPHENYL)-3(R)-(3(S)-HYDROXY-3-((PHENYL OR 4-FLUOROPHENYL))-PROPYL)-4(S)-(4-HYDROXYPHENYL)-2-AZETIDINONE		
(57) Abstract <p>A process for preparing a compound of formula (I) wherein X is H or F, comprising: (a) alkylation of a 3-unsubstituted chiral azetidinone of formula (II) with cinnamyl bromide or 4-fluorocinnamyl bromide; (b) Wacker oxidation of the product of step (a); (c) reduction of the ketone product of step (b); and (d) debenzylation of the ketone of step (c).</p> <div style="text-align: right;"> (I)</div> <div style="text-align: right;"> (II)</div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

5

10 **PROCESS FOR PREPARING 1-(4-FLUOROPHENYL)-3(R)-**
 (3(S)-HYDROXY-3-([PHENYL OR 4-FLUOROPHENYL])-
 PROPYL)-4(S)-(4-HYDROXYPHENYL)-2-AZETIDINONE

15 **BACKGROUND**

 This invention relates to a process for preparing 1-(4-fluorophenyl)-3(R)-(3(S)-hydroxy-3-(4-fluorophenyl)propyl)-4(S)-(4-hydroxyphenyl)-2-azetidinone and 1-(4-fluorophenyl)-3(R)-(3(S)-hydroxy-3-phenylpropyl)-4(S)-(4-hydroxyphenyl)-2-azetidinone,
20 compounds useful as hypocholesterolemic agents, and to the preparation of 1-(4-fluorophenyl)-3(R)-(3-(4-fluorophenyl)-3-oxopropyl)-4(S)-(4-benzyloxyphenyl)-2-azetidinone and 1-(4-fluorophenyl)-3(R)-(3-phenyl-3-oxopropyl)-4(S)-(4-benzyloxyphenyl)-2-azetidinone, intermediates in that process.

25 WO 95/08532 discloses a genus of hypocholesterolemic agents comprising the 3-hydroxypropyl-substituted azetidinones named above and describes several processes suitable for preparing said azetidinones. WO 93/02048 discloses stereoselective processes for producing azetidinones comprising cyclizing intermediates which
30 include the 3-position and 4-position substituents desired in the final product. PCT/US94/07291 discloses a modification of the processes disclosed in WO 93/02048 wherein the β -aminoamides are cyclized by reacting with a silylating agent and a fluoride ion catalyst. WO95/26334 discloses chiral β -amino esters, a process for preparing them, and a
35 process for cyclizing a chiral β -amino ester to obtain a chiral 3-unsubstituted azetidinone. The PCT publications and application are


-2-

related, respectively, to U.S. Serial Nos. 08/257,593 (filed June 9, 1994), 08/178,312 (filed January 11, 1994), 08/089,357 (filed July 9, 1993) and 08/403,081 (filed March 13, 1995), all incorporated herein by reference.

5


SUMMARY OF THE INVENTION

This invention provides a chiral process for preparing a compound having the structural formula I:




10 wherein X is H or F, comprising:

(a) alkylating a chiral 3-unsubstituted azetidinone of formula II with cinnamyl bromide or 4-fluorocinnamyl bromide, III, to obtain a propenyl derivative of formula IV:




15 (b) oxidizing a compound of formula IV to obtain a corresponding ketone of formula V:



(c) reducing a ketone of formula V to a hydroxy compound of formula VI:

-3-



(d) removing the benzyl protecting group from a compound of formula VI to obtain a compound of formula I.

The invention also relates in particular to step (b), wherein
 5 the propenyl intermediate, IV, is oxidized to the ketone, V.

The process of this invention, compared to the procedure for preparing the compounds of formula I disclosed in WO 95/08532, is a simpler procedure, involving fewer steps and milder reagents, and having an improved yield. The ketone intermediate is produced more
 10 efficiently and from a more accessible precursor. The Wacker oxidation step proceeds in a regioselective manner to provide the benzylic oxidation product.

DETAILED DESCRIPTION:

15 In step (a) of the claimed process, the chiral 3-unsubstituted azetidinone of formula II is alkylated with cinnamyl bromide or 4-fluoro-cinnamyl bromide (III) by the method described in WO95/26334, i.e., in an inert anhydrous solvent such as tetrahydrofuran (THF) in the presence of a strong base such as lithium diisopropyl amine
 20 (LDA) at a low temperature of about -70 to -78°C, to obtain a propenyl compound of formula IV.

In step (b), the modified Wacker oxidation comprises treating a propenyl compound of formula IV with an oxidant such as benzoquinone, a catalyst such as palladium acetate, and a catalytic
 25 amount of perchloric acid in a solvent such as acetonitrile:water or dimethylformamide:water at a temperature range of 25-50°C, preferably about 25°C, for 3-12 hours, preferably for about 6 hours. The oxidation product is purified by conventional methods, typically by silica gel chromatography, to obtain the compound of formula V in approximately
 30 a 70% yield. Alternatively, standard Wacker oxidation conditions can be

-4-


used, i.e., palladium chloride and cupric chloride under an oxygen atmosphere.

In step (c), the ketone intermediate of formula V is reduced to the corresponding hydroxy compound, VI, under conditions well known in the art, for example by using a reagent such as NaBH₃ in the presence of a chiral promoting agent such as a chiral borane (e.g., (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrolo-[1,2-c][1,3,2]oxazaborole).

In step (d), the benzyl protecting group on the compound of formula VI is removed by well known methods, for example by hydrogenating an alcoholic solution, e.g., an ethanolic solution, of the compound of formula VI with Pd/C under H₂ at reduced pressure.

The chiral 3-unsubstituted azetidinone starting material of formula II is made by cyclization of a chiral β-amino ester according to the process described in WO95/26334 or the corresponding U.S. application, incorporated herein by reference.

The propenyl compound of formula IV can also be prepared by the process exemplified by the following reaction scheme:




wherein fluorophenyl pentenoic acid, VII, is converted to the corresponding acid chloride and reacted with a chiral auxiliary such as

-5-

- (S)-(+)-4-phenyl-2-oxazolidine. The product, VIII, is enolized and condensed with an imine of formula IX, and the product of that reaction, X, is cyclized by treatment with a silylating agent such as bis trimethylsilylaceta
- 5 tetrabutyl ammonium fluoride (TBAF).


Following are examples of the claimed process.

Example 1




- Cool a solution of 1-(4-fluorophenyl)-4(S)-(4-benzyloxyphenyl)-2-azetidinone (2.5 g, 0.007 mole) in THF (30 mL) to -78°C, add freshly prepared LDA (0.0084 mole) in THF (10 mL) and stir for 1 h at -78°C. Add cinnamyl bromide (1.68 g, 0.0084 mole) dropwise as a THF solution (5 mL). Allow the mixture to warm to -20°C over 4 h and quench with aqueous NH₄Cl (20 mL). Extract the product into ether (50 mL), wash with water (2 X 30 mL) and brine (1 X 30 mL), dry over MgSO₄, filter and concentrate. Purify the resultant crude mixture by flash chromatography on silica gel, eluting with 30% EtOAc/hexanes to obtain 0.268 g (70%) of the title compound plus 0.04 g of the cis isomer. ¹H NMR: (300 MHz, CDCl₃): δ 2.7 (m, 1H), 2.9 (m, 1H), 4.68 (d, J=2 Hz), 5.05 (s, 2H), 6.26 (m, 1H), 6.5 (d, J=12 Hz, 1H), 6.9-7.5 (m, 18H).
- 10
- 15
- 20

Example 2




Step 1:




-6-

Slowly add oxalyl chloride (2.2 g) and DMF (2 drops) to a solution of fluorophenyl pentenoic acid (2.41 g) in CH_2Cl_2 (30 mL) at 0°C , warm the mixture to room temperature and reflux for 1 h. Evaporate the solvent under reduced pressure and azeotrope off the excess oxalyl chloride with CH_2Cl_2 (2 X 50 mL). Redissolve the resultant acid chloride in CH_2Cl_2 (20 mL) and add the solution dropwise to a cooled solution of (S)-(+)-4-phenyl-2-oxazolidone (1.82 g), N, N-diisopropylethylamine (DIPEA) (3.2 g) and N-dimethylaminopyridine (0.056 g) in CH_2Cl_2 (30 mL). Stir the resulting mixture for 10 h, then dilute with CH_2Cl_2 (150 mL). Wash the organic layer with 0.5 N HCl (2 X 50 mL), water (1 X 50 mL) and brine (1 X 50 mL), then dry and concentrate to crystallize the desired product. Recrystallize from ethyl acetate (EtOAc)/hexane and collect the product by filtration to obtain 3.8 g (95%). ^1H NMR: (300 MHz, CDCl_3): δ 2.5 (m, 2H), 3.1 (m, 2H), 4.25 (dd, $J=8$, 3 Hz, 1H), 4.68 (dd, appt. t, $J=8$ Hz), 5.45 (dd, $J=8$, 3 Hz, 1H), 6.1 (m, 1H), 6.35 (d, $J=12$ Hz, 1H), 6.9-7.5 (m, 9H).

Step 2:

Stir a mixture of benzyloxybenzaldehyde (2.12 g, 0.01 mole) and fluoroaniline (1.11 g, 0.01 mole) in isopropanol (25 mL) overnight. Filter to collect the product and dry overnight to obtain 2.8 g.

Step 3:


To a solution of the product of step 1 (1.3 g) in CH_2Cl_2 (20 mL) at -20 to -25°C , slowly add a 1M solution of TiCl_4 in CH_2Cl_2 (3.8 mL) and

-7-

stir at -20°C for 10 min. Slowly add DIPEA (1.3 mL) and stir for 30 min. at -20 °C. Add the product of step 2 (2.34 g) in CH₂Cl₂ (20 mL) and stir for 1 h at a temperature below -20°C. Quench the reaction by adding glacial acetic acid (1.3 mL) in CH₂Cl₂ (3 mL) and stir for 30 min. Pour the mixture into aqueous 2N H₂SO₄ (30 mL), stir for 30 min. and add EtOAc (100 mL). Wash the organic layer with saturated aqueous NaHCO₃ and brine, concentrate and recrystallize from EtOAc/hexane to obtain 1.68 g of the desired product (67%). ¹H NMR: (300 MHz, CDCl₃): δ 2.35 (m, 1H), 2.6 (m, 1H), 4.15 (m, 1H), 4.5 (m, 1H), 4.7 (m, 1H), 5.0 (s, 2H), 5.15 (s, 2H), 5.4 (m, 1H), 6.0 (m, 1H), 6.3 (d, J=12 Hz, 1H), 6.5 (br s, 1H), 6.7-7.9 (m, 22H).

Step 4:

To a suspension of the product of step 3 (1.0 g) in toluene (15 mL) at 60°C, add BSA (1 mL) and heat to 80°C. After 2 h, cool the reaction mixture to 50°C, add TBAF (0.028 g), stir for 2 h more, cool and dilute with EtOAc (100 mL). Wash the organic layer with 0.5 N HCl (2 X 50 mL), water (1 X 50 mL) and brine (1 X 50 mL), dry, filter and concentrate. Purify the crude product by passing through a short pad of silica gel, eluting with 10% EtOAc/hexane to obtain 0.68 g of the title compound (90%). ¹H NMR: (300 MHz, CDCl₃): δ 2.7 (m, 1H), 2.85 (m, 1H), 3.25 (m, 1H), 4.68 (d, J=2Hz, 1H), 5.05 (s, 2H), 6.15 (m, 1H), 6.45 (d, J=12 Hz, 1H), 6.9-7.5 (m, 17H).


Example 3

Combine the product of Example 2 (4.6 mmol), Pd(OAc)₂ (0.0028 g, 2 mol %), benzoquinone (0.07 g, 1.5 eq.) and perchloric acid (catalytic; 0.008 mL) in acetonitrile:water (7:1) (8 mL) and stir for 6 h. Dilute the reaction mixture with EtOAc (50 mL) and wash with water (4 X 25 mL). Dry the organic layer over MgSO₄, filter and treat the filtrate with charcoal (4 g) to remove the palladium catalyst, then filter off the charcoal and concentrate the filtrate. Purify the resultant residue by

-8-


silica gel preparative plate chromatography, eluting with 30 % EtOAc/hexane to obtain the product of step 1 in 70-80% yield. ^1H NMR: (300 MHz, CDCl_3): δ 2.34 (m, 2H), 3.15 (m, 2H), 3.3 (m, 1H), 4.69 (d, $J=1.6$ Hz, 1H), 5.05 (s, 2H), 6.8-7.5 (m, 15H), 8.00 (m, 2H).

5

Example 4

Using the procedure of Example 3, oxidize the azetidinone of Example 1 to obtain the title compound. ^1H NMR: (300 MHz, CDCl_3): δ 2.34 (m, 2H), 3.16 (m, 2H), 3.25 (m, 1H), 4.68 (d, $J=2$ Hz, 1H), 5.04 (s, 2H), 6.9-7.6 (m, 16H), 7.95 (d, 7 Hz, 2H).

10

Example 5

Step 1: Add (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo[1,2-c]-[1,3,2]oxazaborole (120 mg, 0.43 mmol) to 1-(4-fluorophenyl)-3(R)-(3-(4-fluorophenyl)-3-oxopropyl)-4(S)-(4-benzyloxyphenyl)-2-azetidinone (0.95 g, 1.91 mmol) in THF (3 mL) and cool to -20°C . After 5 min., add borohydride-dimethylsulfide complex (2M in THF, 0.85 mL, 1.7 mmol) dropwise over 0.5 h. After a total of 1.5 h, add CH_3OH followed by HCl (1N) and extract the reaction mixture with EtOAc to obtain 1-(4-fluorophenyl)-3(R)-(3(S)-(4-fluorophenyl)-3-hydroxypropyl)-4(S)-(4-benzyloxyphenyl)-2-azetidinone.

15


Step 2: Add 10% Pd/C (0.03 g) to a solution of the product of step 1 (0.4 g, 0.8 mmol) in ethanol (2 mL) and stir under a pressure of H_2 gas (60 psi) for 16 h. Filter the reaction mixture and concentrate to obtain the title compound.

25


-9-

I claim:


1. A process for preparing a compound of the formula



- 5 wherein X is H or F, comprising
(a) alkylation of the 3-unsubstituted azetidinone of the formula




with cinnamyl bromide or 4-fluorocinnamyl bromide to obtain a compound of the formula



10


- (b) oxidation of the product of step (a) to obtain a compound of the formula



- (c) reduction of the ketone of step (b) to obtain a compound of the

15 formula

-10-



; and

(d) debenzylation of the product of step (c).

2. A process of claim 1 comprising:

5 (a) alkylating 1-(4-fluorophenyl)-4(S)-(4-benzyloxyphenyl)-2-azetidinone with cinnamyl bromide or 4-fluorocinnamyl bromide in the presence of a strong base;

(b) oxidizing the product of step (a) with benzoquinone and palladium acetate in the presence of a catalytic amount of perchloric acid;


10 (c) reducing the ketone of step (b) in the presence of a chiral promoting agent; and

(d) debenzylation of the product of step (c) by hydrogenation.

15 3. A process of claim 2 wherein the strong base of step (a) is lithium diisopropyl amine; the reducing agent of step (c) is NaBH_3 and the chiral promoting agent is (R)-tetrahydro-1-methyl-3,3-diphenyl-1H,3H-pyrrolo[1,2-c]-[1,3,2]oxazaborole; and the debenzylation of step (d) comprises hydrogenation of an alcoholic solution of the product of step


20 (c) with palladium on carbon under H_2 at reduced pressure.

4. A process for preparing a compound of the formula



wherein X is H or F, comprising oxidation of a compound of the formula

-11-



wherein X is H or F.

5. A process of claim 4 comprising oxidation with benzoquinone and
5 palladium acetate in the presence of a catalytic amount of perchloric
acid.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/17083

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D205/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 08532 A (SCHERING CORPORATION) 30 March 1995 cited in the application see claims; example 6 ---	1-5
A	WO 95 26334 A (SCHERING CORPORATION) 5 October 1995 cited in the application see claims ---	1-5
A	EP 0 210 705 A (STAMICARBON B.V.) 4 February 1987 -----	

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

23 January 1997

Date of mailing of the international search report

3 1. 01. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat'l Application No
PCT/US 96/17083

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9508532	30-03-95	AU-A- 7795294	10-04-95
		CN-A- 1131416	18-09-96
		CZ-A- 9600839	14-08-96
		EP-A- 0720599	10-07-96
		FI-A- 961300	21-03-96
		HU-A- 73852	30-09-96
		JP-T- 8509989	22-10-96
		NO-A- 961133	20-03-96
		PL-A- 313589	08-07-96
		ZA-A- 9407086	14-03-95
WO-A-9526334	05-10-95	AU-A- 2159695	17-10-95
		CA-A- 2186364	05-10-95
		EP-A- 0751934	08-01-97
		FI-A- 963817	25-09-96
		NO-A- 964008	22-11-96
EP-A-210705	04-02-87	NL-A- 8502144	16-02-87
		JP-A- 62033541	13-02-87